

Saline-immersion peroral endoscopic septotomy: a case report

Roberta Maselli, MD, PhD,^{1,2} Ludovico Alfarone, MD,^{1,2} Davide Massimi, MD,¹ Roberto De Sire, MD,^{1,2} Antonio Capogreco, MD,¹ Alessandro Repici, MD^{1,2}



As an evolution of underwater-assisted endoscopic resection,¹ saline-immersion therapeutic endoscopy has been gaining popularity over the past few years.² Particularly, the saline-immersion technique may become a precious ally for advanced third-space approaches soon. Indeed, the saline-immersion setting has been recently proposed for endoscopic submucosal dissection.^{3,4} Although this assumption is supported by little evidence at the moment, the saline-immersion technique can bring multiple benefits for these complex procedures.

As already well-established for the underwater setting, the saline-immersion technique allows easier scope handling and, thus, better stability. Furthermore, it eliminates any gas–fluid interface within the distal cap, greatly improving the endoscopic view of the submucosal space. Moreover, a better exposure of the submucosal layer is provided by the saline-immersion setting and, hence, a more precise submucosal dissection can be performed. Additionally, saline-aided endotherapy carries significant advantages in terms of safety. The saline-immersion setting makes the detection of the bleeding point easier, which is paramount during the creation of the submucosal tunnel. Furthermore, the replacement of gas with saline avoids gas-related adverse events, such as subcutaneous emphysema and tension capnoperitoneum.

However, beyond these benefits, which are in common with the underwater technique, saline-immersion therapeutic endoscopy may bring some further advantages over underwater methods, especially for third-space approaches. In fact, being a 0.9% saline isotonic solution, as opposed to water, the risk of water intoxication is lower even if a large amount of fluid is instilled in the upper GI tube for long-lasting procedures. Moreover, saline solution allows a better conductivity than the underwater technique, without requiring higher diathermic energy settings for an effective dissection.

Abbreviations: CO₂, carbon dioxide; POES, peroral endoscopic septotomy.

Copyright © 2024 American Society for Gastrointestinal Endoscopy. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

2468-4481

<https://doi.org/10.1016/j.vgie.2023.09.021>

Endoscopy Unit, Department of Gastroenterology, IRCCS Humanitas Research Hospital, Milan, Italy (1), Department of Biomedical Sciences, Humanitas University, Milan, Italy (2).

Here we report the case of a 52-year-old woman with a very short–septum (≤ 20 mm) Zenker diverticulum diagnosed on barium esophagogram and confirmed by endoscopy (Fig. 1). A high-definition gastroscope fitted with a clear distal cap was used. Endotracheal intubation was performed to avoid pulmonary aspiration. We started peroral endoscopic septotomy (POES) under carbon dioxide (CO₂) inflation. First, a submucosal injection was performed on top of the diverticular septum. A mucosal incision was carried out along the long axis of the diverticular septum (Fig. 2). After, a dissection of the immediately underlying submucosa was performed. However, the submucosal space appeared inaccessible because of the small septum. Thus, given the unfeasibility of the procedure under the CO₂ setting, we moved to saline-immersion POES. Thus, CO₂ was switched off, and room temperature sterile saline was infused into the lumen (Fig. 3). After, saline-immersion dissection of the submucosal layer was performed to create 2 short tunnels along both the esophageal and the diverticular sides of the septum (Fig. 4). Then, a complete saline-immersion myotomy was made through the incision of the fully exposed cricopharyngeal muscle fibers (Fig. 5). Finally, complete closure of the mucosal incision site was achieved with 2 hemostatic clips (Fig. 6).

No increase in end-tidal CO₂ or airway pressures was observed during this procedure. No adverse events were reported. After the procedure, the patient remained asymptomatic and afebrile. She was kept fasting for 8 hours and then had a successful trial of clear fluids. The day after the procedure, she resumed oral feeding with a soft diet without any trouble. Over the 2 following years, no recurrence of symptoms was reported by the patient during the scheduled follow-up phone calls.

This pilot experience showed that saline-immersion POES is a feasible option as a rescue therapy for Zenker diverticulum in challenging situations (Video 1, available online at www.videogie.org).

DISCLOSURE

Dr Maselli is a consultant for Fujifilm, ERBE, Apollo Endosurgery, 3D Matrix, and Laborie. Dr Repici is a consultant for Fujifilm, ERBE, Apollo Endosurgery, 3D Matrix, Laborie, and Medtronic. The other authors did not disclose any financial relationships.




Figure 1. Zenker diverticulum with a very small septum.




Figure 2. Mucosal incision along the long axis of the septum.




Figure 3. Saline-immersion access to the submucosal space.



Figure 4. Exposure of the muscular septum through creation of 2 submucosal tunnels.



Figure 5. Saline-immersion myotomy of the muscular septum.




Figure 6. Closure of the mucosal incision site.

REFERENCES

1. Binmoeller KF, Weilert F, Shah J, et al. "Underwater" EMR without submucosal injection for large sessile colorectal polyps (with video). *Gastrointest Endosc* 2012;75:1086-91.
2. Despott EJ, Murino A. Saline-immersion therapeutic endoscopy (SITE): an evolution of underwater endoscopic lesion resection. *Dig Liver Dis* 2017;49:1376.
3. Despott EJ, Hirayama Y, Lazaridis N, et al. Saline immersion therapeutic endoscopy facilitated pocket-creation method for endoscopic submucosal dissection (with video). *Gastrointest Endosc* 2019;89:652-3.
4. Harada H, Nakahara R, Murakami D, et al. Saline-pocket endoscopic submucosal dissection for superficial colorectal neoplasms: a randomized controlled trial (with video). *Gastrointest Endosc* 2019;90:278-87.

Read Articles in Press Online Today!
Visit www.videogie.org

VideoGIE posts in-press articles online in advance of their appearance in a monthly edition of the journal. These articles are available on the *VideoGIE* website by clicking on the "Articles in Press" tab. Articles in Press represent the final edited text of articles that are accepted for publication but not yet scheduled to appear in a specific issue. They are considered officially published as of the date of Web publication, which means readers can access the information and authors can cite the research months prior to its availability in an issue. To cite Articles in Press, include the journal title, year, and the article's Digital Object Identifier (DOI), located in the article footnote. Visit the website today to stay current on the latest research in the field of gastrointestinal endoscopy.